

**REMARKS**

The specification has been amended to correct several errors noted by applicant during a review thereof. The nature of the exact amendments made is apparent from the Appendix hereto in which the changes are shown.

First, the description contains an error with respect to the definition of certain of the rings. These rings were intended to be described as being 4-6 membered saturated rings instead of 5-6 membered saturated rings. The error is believed to be an obvious one because the more specific description and narrow claims name azetidinyI as a possible value of such saturated rings. For example, attention is drawn to to the definition of R<sup>67</sup> on page 14 of the specification. This error has been corrected throughout the specification where appropriate as shown in the Appendix. No new matter is introduced thereby, and entry of these amendments is respectfully requested.

Second, at various locations throughout the specification Examples, the term "mol" was erroneously stated rather than "mmol", and on one occasion the quantity was given as "g" whereas it should have been "mg." Again, both the error and the correction thereof is believed to be readily apparent to a person skilled in this art from like terms used elsewhere throughout the examples. No new matter has been added, and entry of these amendments is respectfully requested.

Third, in several places an unnecessary "or" was included in the specification, and these have been replaced by a ",". These changes also do not constitute new matter, and entry of these amendments is respectfully requested.

Finally, the carbon, hydrogen, nitrogen, oxygen analysis in Example 192 is erroneously given as "C<sub>2</sub>H<sub>1</sub>N<sub>1</sub>O<sub>1</sub>; 1HCL, 2H<sub>2</sub>O", and has been corrected to read

"C<sub>25</sub>H<sub>30</sub>N<sub>4</sub>O<sub>4</sub>". The need for this correction, and the correction to be made, are readily apparent from the compound identified in Example 192 and the percentage analysis given. Again, no new matter has been introduced, and entry of this amendment is respectfully requested.

Original claims 1-30<sup>1</sup> have been cancelled and replaced by new claims 31-60. The replacement claims have been reformatted, so that the metts and bounds of the definitions of the various moieties are easier to read, and have been placed in proper U.S. form by removing "use" language and providing proper dependencies. Claims 31 and 32 have also been substantively amended to remove "direct bond" from the values of Z in claims 1 and 2, so that Z now represents -O-, -NH-, -S- or -CH<sub>2</sub>-. This amendment has been made to further distance the present claims from the disclosures of earlier published PCT applications WO 97/42187 and WO 99/10349, which were cited in the International Search Report and included on the form PTO-1449 submitted with this application as filed.

Original process-for-preparation claims 27-30 referred for their substantive content to various of the Examples. These claims have been cancelled, and new claims 56-60 have been presented, directly reciting process steps based on the Examples as follows:

Claim 56	Example 237
Claim 57	Example 237
Claim 58	Example 237
Claim 59	Example 242
Claim 60	Example 250

<sup>1</sup> Claims 23-30 were added during the International Phase of this PCT application, and a copy of the amended sheet on which they are set out was included with the copy of this application when filed.

Claims 56-60 are therefore fully supported by the respective Examples.

No new matter has been introduced by any of the above amendments, and entry thereof is respectfully requested.


The present application is technically related to an lengthy series of applications filed by applicants' assignee pertaining to to quinazolines and vascular endothelial growth factor receptor tyrosine kinase inhibitors. It is applicants' intent to file a supplemental Information Disclosure Statement citing a potentially vast number of documents which have been cited in one or more of the earlier cases in this series. However, it is presently understood that the references already cited in the International Search Report (and thus in the form PTO-1449 accompanying the U.S. application as filed) are the documents most relevant to the present claims. It is anticipated by the undersigned from past experience with this series that a restriction requirement will be made. Accordingly, it seems most efficient for all concerned to await such restriction requirement and the election of a particular group to prosecute in this application, so that the volume of references cited can be reduced to those most relevant to the elected invention. Moreover, applicants contemplate that further substantive amendments to the claims may be desirable to further distinguish certain embodiments of the presently claimed invention from references already of record (*e.g.*, WO 95/15758), but the ultimate need for, and nature of any such amendment, will likely be affected by a restriction requirement and the particular group elected. Therefore, further substantive amendments are being deferred until it is known whether a restriction requirement will be made and/or the details thereof.

In view of the above-noted further filings being dependant on whether a restriction requirement is made and the nature thereof, and to expedite the prosecution of this

application, it is respectfully requested that the Examiner telephone the undersigned (202-739-5320) if a restriction requirement will not be made in this application, so that appropriate documents and amendments can be filed before a first action on the merits is made. Otherwise, applicants will await the restriction requirement, and file the documents and amendments most appropriate to the elected invention at the time the election is made.

Respectfully Submitted,  
**Morgan Lewis & Bockius LLP**

By: \_\_\_\_\_

  
Donald J. Bird  
Registration No. 25,323  
Tel. No.: (202) 739-5320  
Fax No.: (202) 739-3001

Date: May 6, 2002  
Morgan Lewis & Bockius LLP  
Customer No. **009629**  
1111 Pennsylvania Avenue, N.W.  
Washington, D.C. 20004  
Tel. No.: 202-739-3000  
DJB:mk

**APPENDIX**

**VERSION WITH MARKINGS TO SHOW CHANGES**

**IN THE SPECIFICATION:**

The specification has been amended as shown below, wherein additions are indicated by text in **bold underline**, and deletions are indicated by text **[in bold between brackets]**.

Note that the bolded compound names in the Examples below (which is not also underlined) is bolded in the original specification, and does not represent added text:

The subparagraphs reciting the definitions of R<sup>2</sup> and R<sup>1</sup> from page 3, line 19 to page 9, line 23, have been amended as follows:

R<sup>2</sup> represents hydrogen, hydroxy, halogeno, cyano, nitro, trifluoromethyl, C<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkoxy, C<sub>1-3</sub>alkylsulphanyl, -NR<sup>3</sup>R<sup>4</sup> (wherein R<sup>3</sup> and R<sup>4</sup>, which may be the same or different, each represents hydrogen or C<sub>1-3</sub>alkyl), or R<sup>5</sup>X<sup>1</sup>- (wherein X<sup>1</sup> represents a direct bond, -O-, -CH<sub>2</sub>-, -OC(O)-, -C(O)-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>6</sup>C(O)-, -C(O)NR<sup>7</sup>-, -SO<sub>2</sub>NR<sup>8</sup>-, -NR<sup>9</sup>SO<sub>2</sub>- or -NR<sup>10</sup>- (wherein R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup> and R<sup>10</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl), and R<sup>5</sup> is selected from one of the following twenty-two groups:

- 1) hydrogen, oxiranylC<sub>1-4</sub>alkyl or C<sub>1-5</sub>alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, chloro, bromo and amino;
- 2) C<sub>1-5</sub>alkylX<sup>2</sup>C(O)R<sup>11</sup> (wherein X<sup>2</sup> represents -O- or -NR<sup>12</sup>- (in which R<sup>12</sup> represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>11</sup> represents C<sub>1-3</sub>alkyl, -NR<sup>13</sup>R<sup>14</sup> or -OR<sup>15</sup> (wherein R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup> which may be the same or different each represents hydrogen, C<sub>1-5</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl));
- 3) C<sub>1-5</sub>alkylX<sup>3</sup>R<sup>16</sup> (wherein X<sup>3</sup> represents -O-, -S-, -SO-, -SO<sub>2</sub>-, -OC(O)-, -NR<sup>17</sup>C(O)-, -C(O)NR<sup>18</sup>-, -SO<sub>2</sub>NR<sup>19</sup>-, -NR<sup>20</sup>SO<sub>2</sub>- or -NR<sup>21</sup>- (wherein R<sup>17</sup>, R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and R<sup>21</sup> each

independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>16</sup> represents hydrogen, C<sub>1-3</sub>alkyl, cyclopentyl, cyclohexyl or a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C<sub>1-3</sub>alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C<sub>1-4</sub>alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C<sub>1-4</sub>cyanoalkyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>hydroxyalkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylsulphonylC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxycarbonyl, C<sub>1-4</sub>aminoalkyl, C<sub>1-4</sub>alkylamino, di(C<sub>1-4</sub>alkyl)amino, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkyl, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkoxy, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkoxy and a group  $-(O)_f(C_{1-4}alkyl)_g ringD$  (wherein f is 0 or 1, g is 0 or 1 and ring D is a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C<sub>1-4</sub>alkyl));

4) C<sub>1-5</sub>alkylX<sup>4</sup>C<sub>1-5</sub>alkylX<sup>5</sup>R<sup>22</sup> (wherein X<sup>4</sup> and X<sup>5</sup> which may be the same or different are each -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>23</sup>C(O)-, -C(O)NR<sup>24</sup>-, -SO<sub>2</sub>NR<sup>25</sup>-, -NR<sup>26</sup>SO<sub>2</sub>- or -NR<sup>27</sup>- (wherein R<sup>23</sup>, R<sup>24</sup>, R<sup>25</sup>, R<sup>26</sup> and R<sup>27</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>22</sup> represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl);

5) R<sup>28</sup> (wherein R<sup>28</sup> is a [5]4-6-membered saturated heterocyclic group (linked via carbon or nitrogen) with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C<sub>1-4</sub>cyanoalkyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>hydroxyalkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylsulphonylC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxycarbonyl, C<sub>1-4</sub>aminoalkyl, C<sub>1-4</sub>alkylamino, di(C<sub>1-4</sub>alkyl)amino, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkyl, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkoxy, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkoxy and a group  $-(O)_f(C_{1-4}alkyl)_g ringD$  (wherein f is 0 or 1, g is 0 or 1 and ring D is a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C<sub>1-4</sub>alkyl));

6) C<sub>1-5</sub>alkylR<sup>28</sup> (wherein R<sup>28</sup> is as defined hereinbefore);

7) C<sub>2-5</sub>alkenylR<sup>28</sup> (wherein R<sup>28</sup> is as defined hereinbefore);

8) C<sub>2-5</sub>alkynylR<sup>28</sup> (wherein R<sup>28</sup> is as defined hereinbefore);

- 9)  $R^{29}$  (wherein  $R^{29}$  represents a pyridone group, a phenyl group or a 5-6-membered aromatic heterocyclic group (linked via carbon or nitrogen) with 1-3 heteroatoms selected from O, N and S, which pyridone, phenyl or aromatic heterocyclic group may carry up to 5 substituents selected from hydroxy, halogeno, amino,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ hydroxyalkyl,  $C_{1-4}$ aminoalkyl,  $C_{1-4}$ alkylamino,  $C_{1-4}$ hydroxyalkoxy, carboxy, trifluoromethyl, cyano,  $-C(O)NR^{30}R^{31}$ ,  $-NR^{32}C(O)R^{33}$  (wherein  $R^{30}$ ,  $R^{31}$ ,  $R^{32}$  and  $R^{33}$ , which may be the same or different, each represents hydrogen,  $C_{1-4}$ alkyl or  $C_{1-3}$ alkoxy $C_{2-3}$ alkyl) and a group  $-(O-)_f(C_{1-4}alkyl)_g$ ringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from  $C_{1-4}$ alkyl));
- 10)  $C_{1-5}alkylR^{29}$  (wherein  $R^{29}$  is as defined hereinbefore);
- 11)  $C_{2-5}alkenylR^{29}$  (wherein  $R^{29}$  is as defined hereinbefore);
- 12)  $C_{2-5}alkynylR^{29}$  (wherein  $R^{29}$  is as defined hereinbefore);
- 13)  $C_{1-5}alkylX^6R^{29}$  (wherein  $X^6$  represents  $-O-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-NR^{34}C(O)-$ ,  $-C(O)NR^{35}-$ ,  $-SO_2NR^{36}-$ ,  $-NR^{37}SO_2-$  or  $-NR^{38}-$  (wherein  $R^{34}$ ,  $R^{35}$ ,  $R^{36}$ ,  $R^{37}$  and  $R^{38}$  each independently represents hydrogen,  $C_{1-3}$ alkyl or  $C_{1-3}$ alkoxy $C_{2-3}$ alkyl) and  $R^{29}$  is as defined hereinbefore);
- 14)  $C_{2-5}alkenylX^7R^{29}$  (wherein  $X^7$  represents  $-O-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-NR^{39}C(O)-$ ,  $-C(O)NR^{40}-$ ,  $-SO_2NR^{41}-$ ,  $-NR^{42}SO_2-$  or  $-NR^{43}-$  (wherein  $R^{39}$ ,  $R^{40}$ ,  $R^{41}$ ,  $R^{42}$  and  $R^{43}$  each independently represents hydrogen,  $C_{1-3}$ alkyl or  $C_{1-3}$ alkoxy $C_{2-3}$ alkyl) and  $R^{29}$  is as defined hereinbefore);
- 15)  $C_{2-5}alkynylX^8R^{29}$  (wherein  $X^8$  represents  $-O-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-NR^{44}C(O)-$ ,  $-C(O)NR^{45}-$ ,  $-SO_2NR^{46}-$ ,  $-NR^{47}SO_2-$  or  $-NR^{48}-$  (wherein  $R^{44}$ ,  $R^{45}$ ,  $R^{46}$ ,  $R^{47}$  and  $R^{48}$  each independently represents hydrogen,  $C_{1-3}$ alkyl or  $C_{1-3}$ alkoxy $C_{2-3}$ alkyl) and  $R^{29}$  is as defined hereinbefore);
- 16)  $C_{1-4}alkylX^9C_{1-4}alkylR^{29}$  (wherein  $X^9$  represents  $-O-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-NR^{49}C(O)-$ ,  $-C(O)NR^{50}-$ ,  $-SO_2NR^{51}-$ ,  $-NR^{52}SO_2-$  or  $-NR^{53}-$  (wherein  $R^{49}$ ,  $R^{50}$ ,  $R^{51}$ ,  $R^{52}$  and  $R^{53}$  each independently represents hydrogen,  $C_{1-3}$ alkyl or  $C_{1-3}$ alkoxy $C_{2-3}$ alkyl) and  $R^{29}$  is as defined hereinbefore);
- 17)  $C_{1-4}alkylX^9C_{1-4}alkylR^{28}$  (wherein  $X^9$  and  $R^{28}$  are as defined hereinbefore);
- 18)  $C_{2-5}alkenyl$  which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino,  $C_{1-4}alkylamino$ ,  $N,N$ -di( $C_{1-4}alkyl$ )amino, aminosulphonyl,  $N$ - $C_{1-4}alkylaminosulphonyl$  and  $N,N$ -di( $C_{1-4}alkyl$ )aminosulphonyl;

19) C<sub>2-5</sub>alkynyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, C<sub>1-4</sub>alkylamino, N,N-di(C<sub>1-4</sub>alkyl)amino, aminosulphonyl, N-C<sub>1-4</sub>alkylaminosulphonyl and N,N-di(C<sub>1-4</sub>alkyl)aminosulphonyl;

20) C<sub>2-5</sub>alkenylX<sup>9</sup>C<sub>1-4</sub>alkylR<sup>28</sup> (wherein X<sup>9</sup> and R<sup>28</sup> are as defined hereinbefore);

21) C<sub>2-5</sub>alkynylX<sup>9</sup>C<sub>1-4</sub>alkylR<sup>28</sup> (wherein X<sup>9</sup> and R<sup>28</sup> are as defined hereinbefore); and

22) C<sub>1-4</sub>alkylR<sup>54</sup>(C<sub>1-4</sub>alkyl)<sub>q</sub>(X<sup>9</sup>)<sub>r</sub>R<sup>55</sup> (wherein X<sup>9</sup> is as defined hereinbefore, q is 0 or 1, r is 0 or 1, and R<sup>54</sup> and R<sup>55</sup> are each independently selected from hydrogen, C<sub>1-3</sub>alkyl, cyclopentyl, cyclohexyl and a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C<sub>1-3</sub>alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C<sub>1-4</sub>alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C<sub>1-4</sub>cyanoalkyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>hydroxyalkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylsulphonylC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxycarbonyl, C<sub>1-4</sub>aminoalkyl, C<sub>1-4</sub>alkylamino, di(C<sub>1-4</sub>alkyl)amino, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkyl, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkoxy, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkoxy and a group -(O-)<sub>f</sub>(C<sub>1-4</sub>alkyl)<sub>g</sub>ringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C<sub>1-4</sub>alkyl), with the proviso that R<sup>54</sup> cannot be hydrogen);

and additionally wherein any C<sub>1-5</sub>alkyl, C<sub>2-5</sub>alkenyl or C<sub>2-5</sub>alkynyl group in R<sup>5</sup>X<sup>1</sup>- may bear one or more substituents selected from hydroxy, halogeno and amino);

R<sup>1</sup> represents hydrogen, oxo, halogeno, hydroxy, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxymethyl, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>haloalkyl, cyano, amino, C<sub>2-5</sub>alkenyl, C<sub>2-5</sub>alkynyl, C<sub>1-3</sub>alkanoyloxy, nitro, C<sub>1-4</sub>alkanoylamino, C<sub>1-4</sub>alkoxycarbonyl, C<sub>1-4</sub>alkylsulphonyl, C<sub>1-4</sub>alkylsulphinyl, C<sub>1-4</sub>alkylsulphonyl, carbamoyl, N-C<sub>1-4</sub>alkylcarbamoyl, N,N-di(C<sub>1-4</sub>alkyl)carbamoyl, aminosulphonyl, N-C<sub>1-4</sub>alkylaminosulphonyl, N,N-di(C<sub>1-4</sub>alkyl)aminosulphonyl, N-(C<sub>1-4</sub>alkylsulphonyl)amino, N-(C<sub>1-4</sub>alkylsulphonyl)-N-(C<sub>1-4</sub>alkyl)amino, N,N-di(C<sub>1-4</sub>alkylsulphonyl)amino, a C<sub>3-7</sub>alkylene chain joined to two ring C carbon atoms, C<sub>1-4</sub>alkanoylaminoC<sub>1-4</sub>alkyl, carboxy or a group R<sup>56</sup>X<sup>10</sup> (wherein X<sup>10</sup> represents a direct bond, -O-, -CH<sub>2</sub>-, -OC(O)-, -C(O)-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>57</sup>C(O)-, -C(O)NR<sup>58</sup>-, -SO<sub>2</sub>NR<sup>59</sup>-, -NR<sup>60</sup>SO<sub>2</sub>- or -NR<sup>61</sup>- (wherein R<sup>57</sup>, R<sup>58</sup>, R<sup>59</sup>, R<sup>60</sup> and R<sup>61</sup> each independently represents



hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl), and R<sup>56</sup> is selected from one of the following twenty-two groups:

- 1) hydrogen, oxiranylC<sub>1-4</sub>alkyl or C<sub>1-5</sub>alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, chloro, bromo and amino;
- 2) C<sub>1-5</sub>alkylX<sup>11</sup>C(O)R<sup>62</sup> (wherein X<sup>11</sup> represents -O- or -NR<sup>63</sup>- (in which R<sup>63</sup> represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>62</sup> represents C<sub>1-3</sub>alkyl, -NR<sup>64</sup>R<sup>65</sup> or -OR<sup>66</sup> (wherein R<sup>64</sup>, R<sup>65</sup> and R<sup>66</sup> which may be the same or different each represents hydrogen, C<sub>1-5</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl));
- 3) C<sub>1-5</sub>alkylX<sup>12</sup>R<sup>67</sup> (wherein X<sup>12</sup> represents -O-, -S-, -SO-, -SO<sub>2</sub>-, -OC(O)-, -NR<sup>68</sup>C(O)-, -C(O)NR<sup>69</sup>-, -SO<sub>2</sub>NR<sup>70</sup>-, -NR<sup>71</sup>SO<sub>2</sub>- or -NR<sup>72</sup>- (wherein R<sup>68</sup>, R<sup>69</sup>, R<sup>70</sup>, R<sup>71</sup> and R<sup>72</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>67</sup> represents hydrogen, C<sub>1-3</sub>alkyl, cyclopentyl, cyclohexyl or a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C<sub>1-3</sub>alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C<sub>1-4</sub>alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C<sub>1-4</sub>cyanoalkyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>hydroxyalkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylsulphonylC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxycarbonyl, C<sub>1-4</sub>aminoalkyl, C<sub>1-4</sub>alkylamino, di(C<sub>1-4</sub>alkyl)amino, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkyl, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkoxy, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkoxy and a group -(O-)<sub>f</sub>(C<sub>1-4</sub>alkyl)<sub>g</sub>ringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C<sub>1-4</sub>alkyl));
- 4) C<sub>1-5</sub>alkylX<sup>13</sup>C<sub>1-5</sub>alkylX<sup>14</sup>R<sup>73</sup> (wherein X<sup>13</sup> and X<sup>14</sup> which may be the same or different are each -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>74</sup>C(O)-, -C(O)NR<sup>75</sup>-, -SO<sub>2</sub>NR<sup>76</sup>-, -NR<sup>77</sup>SO<sub>2</sub>- or -NR<sup>78</sup>- (wherein R<sup>74</sup>, R<sup>75</sup>, R<sup>76</sup>, R<sup>77</sup> and R<sup>78</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>73</sup> represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl);
- 5) R<sup>79</sup> (wherein R<sup>79</sup> is a [5]4-6-membered saturated heterocyclic group (linked via carbon or nitrogen) with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C<sub>1-4</sub>cyanoalkyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>hydroxyalkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl, C<sub>1-</sub>

$\text{C}_{1-4}\text{alkylsulphonylC}_{1-4}\text{alkyl}$ ,  $\text{C}_{1-4}\text{alkoxycarbonyl}$ ,  $\text{C}_{1-4}\text{aminoalkyl}$ ,  $\text{C}_{1-4}\text{alkylamino}$ ,  $\text{di}(\text{C}_{1-4}\text{alkyl})\text{amino}$ ,  $\text{C}_{1-4}\text{alkylaminoC}_{1-4}\text{alkyl}$ ,  $\text{di}(\text{C}_{1-4}\text{alkyl})\text{aminoC}_{1-4}\text{alkyl}$ ,  $\text{C}_{1-4}\text{alkylaminoC}_{1-4}\text{alkoxy}$ ,  $\text{di}(\text{C}_{1-4}\text{alkyl})\text{aminoC}_{1-4}\text{alkoxy}$  and a group  $\text{-(O-)}_f(\text{C}_{1-4}\text{alkyl})_g\text{ringD}$  (wherein  $f$  is 0 or 1,  $g$  is 0 or 1 and ring D is a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from  $\text{C}_{1-4}\text{alkyl}$ ));

6)  $\text{C}_{1-5}\text{alkylR}^{79}$  (wherein  $\text{R}^{79}$  is as defined hereinbefore);

7)  $\text{C}_{2-5}\text{alkenylR}^{79}$  (wherein  $\text{R}^{79}$  is as defined hereinbefore);

8)  $\text{C}_{2-5}\text{alkynylR}^{79}$  (wherein  $\text{R}^{79}$  is as defined hereinbefore);

9)  $\text{R}^{80}$  (wherein  $\text{R}^{80}$  represents a pyridone group, a phenyl group or a 5-6-membered aromatic heterocyclic group (linked via carbon or nitrogen) with 1-3 heteroatoms selected from O, N and S, which pyridone, phenyl or aromatic heterocyclic group may carry up to 5 substituents selected from hydroxy, halogeno, amino,  $\text{C}_{1-4}\text{alkyl}$ ,  $\text{C}_{1-4}\text{alkoxy}$ ,  $\text{C}_{1-4}\text{hydroxyalkyl}$ ,  $\text{C}_{1-4}\text{aminoalkyl}$ ,  $\text{C}_{1-4}\text{alkylamino}$ ,  $\text{C}_{1-4}\text{hydroxyalkoxy}$ , carboxy, trifluoromethyl, cyano,  $\text{-C(O)NR}^{81}\text{R}^{82}$ ,  $\text{-NR}^{83}\text{C(O)R}^{84}$  (wherein  $\text{R}^{81}$ ,  $\text{R}^{82}$ ,  $\text{R}^{83}$  and  $\text{R}^{84}$ , which may be the same or different, each represents hydrogen,  $\text{C}_{1-4}\text{alkyl}$  or  $\text{C}_{1-3}\text{alkoxyC}_{2-3}\text{alkyl}$ ) and a group  $\text{-(O-)}_f(\text{C}_{1-4}\text{alkyl})_g\text{ringD}$  (wherein  $f$  is 0 or 1,  $g$  is 0 or 1 and ring D is a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from  $\text{C}_{1-4}\text{alkyl}$ ));

10)  $\text{C}_{1-5}\text{alkylR}^{80}$  (wherein  $\text{R}^{80}$  is as defined hereinbefore);

11)  $\text{C}_{2-5}\text{alkenylR}^{80}$  (wherein  $\text{R}^{80}$  is as defined hereinbefore);

12)  $\text{C}_{2-5}\text{alkynylR}^{80}$  (wherein  $\text{R}^{80}$  is as defined hereinbefore);

13)  $\text{C}_{1-5}\text{alkylX}^{15}\text{R}^{80}$  (wherein  $\text{X}^{15}$  represents  $\text{-O-}$ ,  $\text{-S-}$ ,  $\text{-SO-}$ ,  $\text{-SO}_2\text{-}$ ,  $\text{-NR}^{85}\text{C(O)-}$ ,  $\text{-C(O)NR}^{86}\text{-}$ ,  $\text{-SO}_2\text{NR}^{87}\text{-}$ ,  $\text{-NR}^{88}\text{SO}_2\text{-}$  or  $\text{-NR}^{89}\text{-}$  (wherein  $\text{R}^{85}$ ,  $\text{R}^{86}$ ,  $\text{R}^{87}$ ,  $\text{R}^{88}$  and  $\text{R}^{89}$  each independently represents hydrogen,  $\text{C}_{1-3}\text{alkyl}$  or  $\text{C}_{1-3}\text{alkoxyC}_{2-3}\text{alkyl}$ ) and  $\text{R}^{80}$  is as defined hereinbefore);

14)  $\text{C}_{2-5}\text{alkenylX}^{16}\text{R}^{80}$  (wherein  $\text{X}^{16}$  represents  $\text{-O-}$ ,  $\text{-S-}$ ,  $\text{-SO-}$ ,  $\text{-SO}_2\text{-}$ ,  $\text{-NR}^{90}\text{C(O)-}$ ,  $\text{-C(O)NR}^{91}\text{-}$ ,  $\text{-SO}_2\text{NR}^{92}\text{-}$ ,  $\text{-NR}^{93}\text{SO}_2\text{-}$  or  $\text{-NR}^{94}\text{-}$  (wherein  $\text{R}^{90}$ ,  $\text{R}^{91}$ ,  $\text{R}^{92}$ ,  $\text{R}^{93}$  and  $\text{R}^{94}$  each independently represents hydrogen,  $\text{C}_{1-3}\text{alkyl}$  or  $\text{C}_{1-3}\text{alkoxyC}_{2-3}\text{alkyl}$ ) and  $\text{R}^{80}$  is as defined hereinbefore);

- 15)  $C_{2-5}alkynylX^{17}R^{80}$  (wherein  $X^{17}$  represents -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>95</sup>C(O)-, -C(O)NR<sup>96</sup>-, -SO<sub>2</sub>NR<sup>97</sup>-, -NR<sup>98</sup>SO<sub>2</sub>- or -NR<sup>99</sup>- (wherein R<sup>95</sup>, R<sup>96</sup>, R<sup>97</sup>, R<sup>98</sup> and R<sup>99</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>80</sup> is as defined hereinbefore);
- 16)  $C_{1-4}alkylX^{18}C_{1-4}alkylR^{80}$  (wherein  $X^{18}$  represents -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>100</sup>C(O)-, -C(O)NR<sup>101</sup>-, -SO<sub>2</sub>NR<sup>102</sup>-, -NR<sup>103</sup>SO<sub>2</sub>- or -NR<sup>104</sup>- (wherein R<sup>100</sup>, R<sup>101</sup>, R<sup>102</sup>, R<sup>103</sup> and R<sup>104</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>80</sup> is as defined hereinbefore);
- 17)  $C_{1-4}alkylX^{18}C_{1-4}alkylR^{79}$  (wherein  $X^{18}$  and R<sup>79</sup> are as defined hereinbefore);
- 18) C<sub>2-5</sub>alkenyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, C<sub>1-4</sub>alkylamino, N,N-di(C<sub>1-4</sub>alkyl)amino, aminosulphonyl, N-C<sub>1-4</sub>alkylaminosulphonyl and N,N-di(C<sub>1-4</sub>alkyl)aminosulphonyl;
- 19) C<sub>2-5</sub>alkynyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, C<sub>1-4</sub>alkylamino, N,N-di(C<sub>1-4</sub>alkyl)amino, aminosulphonyl, N-C<sub>1-4</sub>alkylaminosulphonyl and N,N-di(C<sub>1-4</sub>alkyl)aminosulphonyl;
- 20)  $C_{2-5}alkenylX^{18}C_{1-4}alkylR^{79}$  (wherein  $X^{18}$  and R<sup>79</sup> are as defined hereinbefore);
- 21)  $C_{2-5}alkynylX^{18}C_{1-4}alkylR^{79}$  (wherein  $X^{18}$  and R<sup>79</sup> are as defined hereinbefore); and
- 22)  $C_{1-4}alkylR^{105}(C_{1-4}alkyl)_x(X^{18})_yR^{106}$  (wherein  $X^{18}$  is as defined hereinbefore, x is 0 or 1, y is 0 or 1, and R<sup>105</sup> and R<sup>106</sup> are each independently selected from hydrogen, C<sub>1-3</sub>alkyl, cyclopentyl, cyclohexyl and a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C<sub>1-3</sub>alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C<sub>1-4</sub>alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C<sub>1-4</sub>cyanoalkyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>hydroxyalkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylsulphonylC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxycarbonyl, C<sub>1-4</sub>aminoalkyl, C<sub>1-4</sub>alkylamino, di(C<sub>1-4</sub>alkyl)amino, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkyl, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkoxy, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkoxy and a group -(O-)<sub>f</sub>(C<sub>1-4</sub>alkyl)<sub>g</sub>ringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C<sub>1-4</sub>alkyl) with the proviso that R<sup>105</sup> cannot be hydrogen);

and additionally wherein any C<sub>1-5</sub>alkyl, C<sub>2-5</sub>alkenyl or C<sub>2-5</sub>alkynyl group in R<sup>56</sup>X<sup>10</sup>- may bear one or more substituents selected from hydroxy, halogeno and amino);  
or a salt thereof, or a prodrug thereof for example an ester or an amide, in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in warm-blooded animals such as humans.

\* \* \* \* \*

The paragraph on page 15, extending from line 24 to line 32 has been amended as follows:

Advantageously R<sup>105</sup> and R<sup>106</sup> are each independently a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C<sub>1-3</sub>cyanoalkyl, C<sub>1-3</sub>alkyl, C<sub>1-3</sub>hydroxyalkyl, C<sub>1-3</sub>alkoxy, C<sub>1-2</sub>alkoxyC<sub>1-3</sub>alkyl, C<sub>1-2</sub>alkylsulphonylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkoxycarbonyl, C<sub>1-3</sub>alkylamino, di(C<sub>1-3</sub>alkyl)amino, C<sub>1-3</sub>alkylaminoC<sub>1-3</sub>alkyl, di(C<sub>1-3</sub>alkyl)aminoC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkylaminoC<sub>1-3</sub>alkoxy, di(C<sub>1-3</sub>alkyl)aminoC<sub>1-3</sub>alkoxy and a group  $-(O)-(C_{1-3}alkyl)_g ringD$  (wherein f is 0 or 1, g is 0 or 1 and ring D is a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C<sub>1-3</sub>alkyl).

\* \* \* \* \*

The paragraph extending from page 20, line 30 to page 21, line 5 has been amended as follows:

Advantageously R<sup>54</sup> and R<sup>55</sup> are each independently a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C<sub>1-3</sub>cyanoalkyl, C<sub>1-3</sub>alkyl, C<sub>1-3</sub>hydroxyalkyl, C<sub>1-3</sub>alkoxy, C<sub>1-2</sub>alkoxyC<sub>1-3</sub>alkyl, C<sub>1-2</sub>alkylsulphonylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkoxycarbonyl and a group  $-(O)-(C_{1-3}alkyl)_g ringD$  (wherein f is 0 or 1, g is 0 or 1 and ring D is a [5]4-6-membered saturated heterocyclic group with 1-2

heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C<sub>1-3</sub>alkyl).

\* \* \* \* \*

The paragraphs extending from page 22, line 7 to page 23, line 8 has been amended as follows:

3) C<sub>2-4</sub>alkylX<sup>3</sup>R<sup>16</sup> (wherein X<sup>3</sup> is as hereinbefore defined and R<sup>16</sup> represents hydrogen, C<sub>1-3</sub>alkyl, cyclopentyl, cyclohexyl or a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C<sub>1-3</sub>alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C<sub>1-3</sub>alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C<sub>1-4</sub>cyanoalkyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>hydroxyalkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylsulphonylC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxycarbonyl, C<sub>1-4</sub>alkylamino, di(C<sub>1-4</sub>alkyl)amino, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkyl, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkoxy, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkoxy and a group -(-O-)<sub>f</sub>(C<sub>1-4</sub>alkyl)<sub>g</sub>ringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C<sub>1-4</sub>alkyl));

4) C<sub>2-3</sub>alkylX<sup>4</sup>C<sub>2-3</sub>alkylX<sup>5</sup>R<sup>22</sup> (wherein X<sup>4</sup> and X<sup>5</sup> are as hereinbefore defined and R<sup>22</sup> represents hydrogen or C<sub>1-3</sub>alkyl);

5) R<sup>28</sup> (wherein R<sup>28</sup> is as defined hereinbefore);

6) C<sub>1-5</sub>alkylR<sup>107</sup> (wherein R<sup>107</sup> is a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group is linked to C<sub>1-5</sub>alkyl through a carbon atom and which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C<sub>1-4</sub>cyanoalkyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>hydroxyalkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylsulphonylC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxycarbonyl, C<sub>1-4</sub>alkylamino, di(C<sub>1-4</sub>alkyl)amino, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkyl, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkoxy, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkoxy and a group -(-O-)<sub>f</sub>(C<sub>1-4</sub>alkyl)<sub>g</sub>ringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C<sub>1-4</sub>alkyl)) or C<sub>2-5</sub>alkylR<sup>108</sup> (wherein R<sup>108</sup> is a [5]4-6-

membered saturated heterocyclic group with 1-2 heteroatoms, of which one is N and the other may be selected independently from O, S and N, which heterocyclic group is linked to C<sub>2-5</sub>alkyl through a nitrogen atom and which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C<sub>1-4</sub>cyanoalkyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>hydroxyalkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylsulphonylC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxycarbonyl, C<sub>1-4</sub>alkylamino, di(C<sub>1-4</sub>alkyl)amino, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkyl, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkoxy, di(C<sub>1-4</sub>alkyl)aminoC<sub>1-4</sub>alkoxy and a group  $-(O-)_f(C_{1-4}alkyl)_{g}ringD$  (wherein f is 0 or 1, g is 0 or 1 and ring D is a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C<sub>1-4</sub>alkyl));

\* \* \* \* \*

The paragraph extending from page 28, line 12 to page 30, line 21 has been amended as follows:

More preferably R<sup>2</sup> represents hydroxy, C<sub>1-3</sub>alkyl, amino or R<sup>5</sup>X<sup>1</sup> - [wherein X<sup>1</sup> is as hereinbefore defined and R<sup>5</sup> represents methyl, ethyl, benzyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-methoxyethyl, 3-methoxypropyl, 2-(methylsulphinyl)ethyl, 2-(methylsulphonyl)ethyl, 2-(ethylsulphinyl)ethyl, 2-(ethylsulphonyl)ethyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl, 2-sulphamoyl ethyl, 2-(methylamino)ethyl, 3-(methylamino)propyl, 2-(ethylamino)ethyl, 3-(ethylamino)propyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-(N,N-diethylamino)ethyl, 3-(N,N-diethylamino)propyl, 2-(N-methyl-N-methylsulphonylamino)ethyl, 3-(N-methyl-N-methylsulphonylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(methylpiperidino)ethyl, 3-(methylpiperidino)propyl, 2-(ethylpiperidino)ethyl, 3-(ethylpiperidino)propyl, 2-((2-methoxyethyl)piperidino)ethyl, 3-((2-methoxyethyl)piperidino)propyl, 2-((2-methylsulphonyl)ethylpiperidino)ethyl, 3-((2-methylsulphonyl)ethylpiperidino)propyl, piperidin-3-ylmethyl, piperidin-4-ylmethyl, 2-(piperidin-3-yl)ethyl, 2-(piperidin-4-yl)ethyl, 3-(piperidin-3-yl)propyl, 3-(piperidin-4-yl)propyl, 2-(piperidin-2-yl)ethyl, 3-(piperidin-2-yl)propyl, (1-methylpiperidin-3-yl)methyl,

(1-methylpiperidin-4-yl)methyl, (1-cyanomethylpiperidin-3-yl)methyl, (1-cyanomethylpiperidin-4-yl)methyl, 2-(methylpiperidin-3-yl)ethyl, 2-(methylpiperidin-4-yl)ethyl, 2-(1-cyanomethylpiperidin-3-yl)ethyl, 2-(1-cyanomethylpiperidin-4-yl)ethyl, 3-(methylpiperidin-3-yl)propyl, 3-(methylpiperidin-4-yl)propyl, 3-(1-cyanomethylpiperidin-3-yl)propyl, 3-(1-cyanomethylpiperidin-4-yl)propyl, 2-(ethylpiperidin-3-yl)ethyl, 2-(ethylpiperidin-4-yl)ethyl, 3-(ethylpiperidin-3-yl)propyl, 3-(ethylpiperidin-4-yl)propyl, ((2-methoxyethyl)piperidin-3-yl)methyl, ((2-methoxyethyl)piperidin-4-yl)methyl, 2-((2-methoxyethyl)piperidin-3-yl)ethyl, 2-((2-methoxyethyl)piperidin-4-yl)ethyl, 3-((2-methoxyethyl)piperidin-3-yl)propyl, 3-((2-methoxyethyl)piperidin-4-yl)propyl, (1-(2-methylsulphonyl)ethyl)piperidin-3-yl)methyl, (1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methyl, 2-((2-methylsulphonyl)ethyl)piperidin-3-yl)ethyl, 2-((2-methylsulphonyl)ethyl)piperidin-4-yl)ethyl, 3-((2-methylsulphonyl)ethyl)piperidin-3-yl)propyl, 3-((2-methylsulphonyl)ethyl)piperidin-4-yl)propyl, 1-isopropylpiperidin-2-yl)methyl, 1-isopropylpiperidin-3-yl)methyl, 1-isopropylpiperidin-4-yl)methyl, 2-(1-isopropylpiperidin-2-yl)ethyl, 2-(1-isopropylpiperidin-3-yl)ethyl, 2-(1-isopropylpiperidin-4-yl)ethyl, 3-(1-isopropylpiperidin-2-yl)propyl, 3-(1-isopropylpiperidin-3-yl)propyl, 3-(1-isopropylpiperidin-4-yl)propyl, 2-(piperidin-4-yloxy)ethyl, 3-(piperidin-4-yloxy)propyl, 2-(1-(cyanomethyl)piperidin-4-yloxy)ethyl, 3-(1-(cyanomethyl)piperidin-4-yloxy)propyl, 2-(1-(2-cyanoethyl)piperidin-4-yloxy)ethyl, 3-(1-(2-cyanoethyl)piperidin-4-yloxy)propyl, 2-(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, (pyrrolidin-2-yl)methyl, 2-(pyrrolidin-1-yl)ethyl, 3-(pyrrolidin-1-yl)propyl, (2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, 5(R)-(2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, (5S)-(2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(N-(2-methoxyethyl)-N-methylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-(2-methoxyethylamino)propyl, 3-(N-(2-methoxyethyl)-N-methylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-methylthiazol-4-yl)methyl, 2-acetamidothiazol-4-yl)methyl, 1-methylimidazol-2-yl)methyl, 2-(imidazol-1-yl)ethyl, 2-(2-methylimidazol-1-yl)ethyl, 2-(2-ethylimidazol-1-yl)ethyl, 3-(2-methylimidazol-1-yl)propyl, 3-(2-ethylimidazol-1-yl)propyl, 2-(1,2,3-triazol-1-yl)ethyl, 2-(1,2,3-triazol-2-yl)ethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-(1,2,4-triazol-4-yl)ethyl, 4-pyridyl)methyl, 2-(4-pyridyl)ethyl, 3-(4-

pyridyl)propyl, 2-(4-pyridyloxy)ethyl, 2-(4-pyridylamino)ethyl, 2-(4-oxo-1,4-dihydro-1-pyridyl)ethyl, 2-(2-oxo-imidazolidin-1-yl)ethyl, 3-(2-oxo-imidazolidin-1-yl)propyl, 2-thiomorpholinoethyl, 3-thiomorpholinopropyl, 2-(1,1-dioxothiomorpholino)ethyl, 3-(1,1-dioxothiomorpholino)propyl, 2-(2-methoxyethoxy)ethyl, 2-(4-methylpiperazin-1-yl)ethyl, 3-(4-methylpiperazin-1-yl)propyl, 3-(methylsulphinyl)propyl, 3-(methylsulphonyl)propyl, 3-(ethylsulphinyl)propyl, 3-(ethylsulphonyl)propyl, 2-(5-methyl-1,2,4-triazol-1-yl)ethyl, morpholino, 2-((N-(1-methylimidazol-4-ylsulphonyl)-N-methyl)amino)ethyl, 2-((N-(3-morpholinopropylsulphonyl)-N-methyl)amino)ethyl, 2-((N-methyl-N-4-pyridyl)amino)ethyl, 3-(4-oxidomorpholino)propyl, 2-(2-(4-methylpiperazin-1-yl)ethoxy)ethyl, 3-(2-(4-methylpiperazin-1-yl)ethoxy)propyl, 2-(2-morpholinoethoxy)ethyl, 3-(2-morpholinoethoxy)propyl, 2-(tetrahydropyran-4-yloxy)ethyl, 3-(tetrahydropyran-4-yloxy)propyl, 2-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)vinyl, 3-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)prop-2-en-1-yl, 1-(2-pyrrolidinylethyl)piperidin-4-ylmethyl, 1-(3-pyrrolidinylpropyl)piperidin-4-ylmethyl, 1-(2-piperidinylethyl)piperidin-4-ylmethyl, 1-(3-piperidinylpropyl)piperidin-4-ylmethyl, 1-(2-morpholinoethyl)piperidin-4-ylmethyl, 1-(3-morpholinopropyl)piperidin-4-ylmethyl, 1-(2-thiomorpholinoethyl)piperidin-4-ylmethyl, 1-(3-thiomorpholinopropyl)piperidin-4-ylmethyl, 1-(2-azetidinyethyl)piperidin-4-ylmethyl, [or] 1-(3-azetidinypropyl)piperidin-4-ylmethyl, 3-morpholino-2-hydroxypropyl, (2*R*)-3-morpholino-2-hydroxypropyl, (2*S*)-3-morpholino-2-hydroxypropyl, 3-piperidino-2-hydroxypropyl, (2*R*)-3-piperidino-2-hydroxypropyl, (2*S*)-3-piperidino-2-hydroxypropyl, 3-pyrrolidin-1-yl-2-hydroxypropyl, (2*R*)-3-pyrrolidin-1-yl-2-hydroxypropyl, (2*S*)-3-pyrrolidin-1-yl-2-hydroxypropyl, 3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, (2*R*)-3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, (2*S*)-3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, 3-(N,N-diethylamino)-2-hydroxypropyl, (2*R*)-3-(N,N-diethylamino)-2-hydroxypropyl, (2*S*)-3-(N,N-diethylamino)-2-hydroxypropyl, 3-(isopropylamino)-2-hydroxypropyl, (2*R*)-3-(isopropylamino)-2-hydroxypropyl, (2*S*)-3-(isopropylamino)-2-hydroxypropyl, 3-(N,N-diisopropylamino)-2-hydroxypropyl, (2*R*)-3-(N,N-diisopropylamino)-2-hydroxypropyl or (2*S*)-3-(N,N-diisopropylamino)-2-hydroxypropyl].

\* \* \* \* \*



The paragraph extending from page 30, line 22 to page 32, line 30 has been amended as follows:

Particularly  $R^2$  represents  $C_{1-3}$ alkyl, amino or  $R^5X^1$  - [wherein  $X^1$  is as hereinbefore defined and  $R^5$  represents ethyl, benzyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-methoxyethyl, 3-methoxypropyl, 2-(methylsulphinyl)ethyl, 2-(methylsulphonyl)ethyl, 2-(ethylsulphinyl)ethyl, 2-(ethylsulphonyl)ethyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl, 2-sulphamoylethyl, 2-(methylamino)ethyl, 3-(methylamino)propyl, 2-(ethylamino)ethyl, 3-(ethylamino)propyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-(N,N-diethylamino)ethyl, 3-(N,N-diethylamino)propyl, 2-(N-methyl-N-methylsulphonylamino)ethyl, 3-(N-methyl-N-methylsulphonylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(methylpiperidino)ethyl, 3-(methylpiperidino)propyl, 2-(ethylpiperidino)ethyl, 3-(ethylpiperidino)propyl, 2-((2-methoxyethyl)piperidino)ethyl, 3-((2-methoxyethyl)piperidino)propyl, 2-((2-methylsulphonyl)ethylpiperidino)ethyl, 3-((2-methylsulphonyl)ethylpiperidino)propyl, piperidin-3-ylmethyl, piperidin-4-ylmethyl, 2-(piperidin-3-yl)ethyl, 2-(piperidin-4-yl)ethyl, 3-(piperidin-3-yl)propyl, 3-(piperidin-4-yl)propyl, 2-(piperidin-2-yl)ethyl, 3-(piperidin-2-yl)propyl, (1-methylpiperidin-3-yl)methyl, (1-methylpiperidin-4-yl)methyl, (1-cyanomethylpiperidin-3-yl)methyl, (1-cyanomethylpiperidin-4-yl)methyl, 2-(methylpiperidin-3-yl)ethyl, 2-(methylpiperidin-4-yl)ethyl, 2-(1-cyanomethylpiperidin-3-yl)ethyl, 2-(1-cyanomethylpiperidin-4-yl)ethyl, 3-(methylpiperidin-3-yl)propyl, 3-(methylpiperidin-4-yl)propyl, 3-(1-cyanomethylpiperidin-3-yl)propyl, 3-(1-cyanomethylpiperidin-4-yl)propyl, 2-(ethylpiperidin-3-yl)ethyl, 2-(ethylpiperidin-4-yl)ethyl, 3-(ethylpiperidin-3-yl)propyl, 3-(ethylpiperidin-4-yl)propyl, ((2-methoxyethyl)piperidin-3-yl)methyl, ((2-methoxyethyl)piperidin-4-yl)methyl, 2-((2-methoxyethyl)piperidin-3-yl)ethyl, 2-((2-methoxyethyl)piperidin-4-yl)ethyl, 3-((2-methoxyethyl)piperidin-3-yl)propyl, 3-((2-methoxyethyl)piperidin-4-yl)propyl, (1-(2-methylsulphonylethyl)piperidin-3-yl)methyl, (1-(2-methylsulphonylethyl)piperidin-4-yl)methyl, 2-((2-methylsulphonylethyl)piperidin-3-yl)ethyl, 2-((2-methylsulphonylethyl)piperidin-4-yl)ethyl, 3-((2-methylsulphonylethyl)piperidin-3-yl)propyl, 3-((2-methylsulphonylethyl)piperidin-4-yl)propyl, 1-isopropylpiperidin-2-ylmethyl, 1-

isopropylpiperidin-3-ylmethyl, 1-isopropylpiperidin-4-ylmethyl, 2-(1-isopropylpiperidin-2-yl)ethyl, 2-(1-isopropylpiperidin-3-yl)ethyl, 2-(1-isopropylpiperidin-4-yl)ethyl, 3-(1-isopropylpiperidin-2-yl)propyl, 3-(1-isopropylpiperidin-3-yl)propyl, 3-(1-isopropylpiperidin-4-yl)propyl, 2-(piperidin-4-yloxy)ethyl, 3-(piperidin-4-yloxy)propyl, 2-(1-(cyanomethyl)piperidin-4-yloxy)ethyl, 3-(1-(cyanomethyl)piperidin-4-yloxy)propyl, 2-(1-(2-cyanoethyl)piperidin-4-yloxy)ethyl, 3-(1-(2-cyanoethyl)piperidin-4-yloxy)propyl, 2-(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, (pyrrolidin-2-yl)methyl, 2-(pyrrolidin-1-yl)ethyl, 3-(pyrrolidin-1-yl)propyl, (2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, 5(R)-(2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, (5S)-(2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(N-(2-methoxyethyl)-N-methylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-(2-methoxyethylamino)propyl, 3-(N-(2-methoxyethyl)-N-methylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-methylthiazol-4-ylmethyl, 2-acetamidothiazol-4-ylmethyl, 1-methylimidazol-2-ylmethyl, 2-(imidazol-1-yl)ethyl, 2-(2-methylimidazol-1-yl)ethyl, 2-(2-ethylimidazol-1-yl)ethyl, 3-(2-methylimidazol-1-yl)propyl, 3-(2-ethylimidazol-1-yl)propyl, 2-(1,2,3-triazol-1-yl)ethyl, 2-(1,2,3-triazol-2-yl)ethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-(1,2,4-triazol-4-yl)ethyl, 4-pyridylmethyl, 2-(4-pyridyl)ethyl, 3-(4-pyridyl)propyl, 2-(4-pyridyloxy)ethyl, 2-(4-pyridylamino)ethyl, 2-(4-oxo-1,4-dihydro-1-pyridyl)ethyl, 2-(2-oxo-imidazolidin-1-yl)ethyl, 3-(2-oxo-imidazolidin-1-yl)propyl, 2-thiomorpholinoethyl, 3-thiomorpholinopropyl, 2-(1,1-dioxothiomorpholino)ethyl, 3-(1,1-dioxothiomorpholino)propyl, 2-(2-methoxyethoxy)ethyl, 2-(4-methylpiperazin-1-yl)ethyl, 3-(4-methylpiperazin-1-yl)propyl, 3-(methylsulphinyl)propyl, 3-(methylsulphonyl)propyl, 3-(ethylsulphinyl)propyl, 3-(ethylsulphonyl)propyl, 2-(5-methyl-1,2,4-triazol-1-yl)ethyl, morpholino, 2-(N-(1-methylimidazol-4-ylsulphonyl)-N-methyl)amino)ethyl, 2-(N-(3-morpholinopropylsulphonyl)-N-methyl)amino)ethyl, 2-(N-methyl-N-4-pyridyl)amino)ethyl, 3-(4-oxidomorpholino)propyl, 2-(2-(4-methylpiperazin-1-yl)ethoxy)ethyl, 3-(2-(4-methylpiperazin-1-yl)ethoxy)propyl, 2-(2-morpholinoethoxy)ethyl, 3-(2-morpholinoethoxy)propyl, 2-(tetrahydropyran-4-yloxy)ethyl, 3-(tetrahydropyran-4-yloxy)propyl, 2-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)vinyl, 3-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)prop-2-en-1-yl, 1-(2-pyrrolidinylethyl)piperidin-4-ylmethyl, 1-(3-pyrrolidinylpropyl)piperidin-4-ylmethyl, 1-(2-piperidinylethyl)piperidin-4-ylmethyl, 1-

(3-piperidinylpropyl)piperidin-4-ylmethyl, 1-(2-morpholinoethyl)piperidin-4-ylmethyl, 1-(3-morpholinopropyl)piperidin-4-ylmethyl, 1-(2-thiomorpholinoethyl)piperidin-4-ylmethyl, 1-(3-thiomorpholinopropyl)piperidin-4-ylmethyl, 1-(2-azetidinyloethyl)piperidin-4-ylmethyl, [or] 1-(3-azetidinylopropyl)piperidin-4-ylmethyl, 3-morpholino-2-hydroxypropyl, (2R)-3-morpholino-2-hydroxypropyl, (2S)-3-morpholino-2-hydroxypropyl, 3-piperidino-2-hydroxypropyl, (2R)-3-piperidino-2-hydroxypropyl, (2S)-3-piperidino-2-hydroxypropyl, 3-pyrrolidin-1-yl-2-hydroxypropyl, (2R)-3-pyrrolidin-1-yl-2-hydroxypropyl, (2S)-3-pyrrolidin-1-yl-2-hydroxypropyl, 3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, (2R)-3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, (2S)-3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, 3-(N,N-diethylamino)-2-hydroxypropyl, (2R)-3-(N,N-diethylamino)-2-hydroxypropyl, (2S)-3-(N,N-diethylamino)-2-hydroxypropyl, 3-(isopropylamino)-2-hydroxypropyl, (2R)-3-(isopropylamino)-2-hydroxypropyl, (2S)-3-(isopropylamino)-2-hydroxypropyl, 3-(N,N-diisopropylamino)-2-hydroxypropyl, (2R)-3-(N,N-diisopropylamino)-2-hydroxypropyl or (2S)-3-(N,N-diisopropylamino)-2-hydroxypropyl].

\* \* \* \* \*

The paragraph extending from page 32, line 31 to page 35, line 3 has been amended as follows:

More particularly R<sup>2</sup> represents C<sub>1-3</sub>alkyl, amino or R<sup>5</sup>X<sup>1</sup> - [wherein X<sup>1</sup> is as hereinbefore defined and R<sup>5</sup> represents ethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-methoxyethyl, 3-methoxypropyl, 2-(methylsulphinyl)ethyl, 2-(methylsulphonyl)ethyl, 2-(ethylsulphinyl)ethyl, 2-(ethylsulphonyl)ethyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl, 2-sulphamoyl, 2-(methylamino)ethyl, 3-(methylamino)propyl, 2-(ethylamino)ethyl, 3-(ethylamino)propyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-(N,N-diethylamino)ethyl, 3-(N,N-diethylamino)propyl, 2-(N-methyl-N-methylsulphonylamino)ethyl, 3-(N-methyl-N-methylsulphonylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(methylpiperidino)ethyl, 3-(methylpiperidino)propyl, 2-(ethylpiperidino)ethyl, 3-(ethylpiperidino)propyl, 2-((2-methoxyethyl)piperidino)ethyl, 3-((2-methoxyethyl)piperidino)propyl, 2-((2-methylsulphonyl)ethylpiperidino)ethyl, 3-((2-

methylsulphonyl)ethylpiperidino)propyl, piperidin-3-ylmethyl, piperidin-4-ylmethyl, 2-(piperidin-3-yl)ethyl, 2-(piperidin-4-yl)ethyl, 3-(piperidin-3-yl)propyl, 3-(piperidin-4-yl)propyl, 2-(piperidin-2-yl)ethyl, 3-(piperidin-2-yl)propyl, (1-methylpiperidin-3-yl)methyl, (1-methylpiperidin-4-yl)methyl, (1-cyanomethylpiperidin-3-yl)methyl, (1-cyanomethylpiperidin-4-yl)methyl, 2-(methylpiperidin-3-yl)ethyl, 2-(methylpiperidin-4-yl)ethyl, 2-(1-cyanomethylpiperidin-3-yl)ethyl, 2-(1-cyanomethylpiperidin-4-yl)ethyl, 3-(methylpiperidin-3-yl)propyl, 3-(methylpiperidin-4-yl)propyl, 3-(1-cyanomethylpiperidin-3-yl)propyl, 3-(1-cyanomethylpiperidin-4-yl)propyl, 2-(ethylpiperidin-3-yl)ethyl, 2-(ethylpiperidin-4-yl)ethyl, 3-(ethylpiperidin-3-yl)propyl, 3-(ethylpiperidin-4-yl)propyl, ((2-methoxyethyl)piperidin-3-yl)methyl, ((2-methoxyethyl)piperidin-4-yl)methyl, 2-((2-methoxyethyl)piperidin-3-yl)ethyl, 2-((2-methoxyethyl)piperidin-4-yl)ethyl, 3-((2-methoxyethyl)piperidin-3-yl)propyl, 3-((2-methoxyethyl)piperidin-4-yl)propyl, (1-(2-methylsulphonyl)ethyl)piperidin-3-yl)methyl, (1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methyl, 2-((2-methylsulphonyl)ethyl)piperidin-3-yl)ethyl, 2-((2-methylsulphonyl)ethyl)piperidin-4-yl)ethyl, 3-((2-methylsulphonyl)ethyl)piperidin-3-yl)propyl, 3-((2-methylsulphonyl)ethyl)piperidin-4-yl)propyl, 1-isopropylpiperidin-2-ylmethyl, 1-isopropylpiperidin-3-ylmethyl, 1-isopropylpiperidin-4-ylmethyl, 2-(1-isopropylpiperidin-2-yl)ethyl, 2-(1-isopropylpiperidin-3-yl)ethyl, 2-(1-isopropylpiperidin-4-yl)ethyl, 3-(1-isopropylpiperidin-2-yl)propyl, 3-(1-isopropylpiperidin-3-yl)propyl, 3-(1-isopropylpiperidin-4-yl)propyl, 2-(piperidin-4-yloxy)ethyl, 3-(piperidin-4-yloxy)propyl, 2-(1-(cyanomethyl)piperidin-4-yloxy)ethyl, 3-(1-(cyanomethyl)piperidin-4-yloxy)propyl, 2-(1-(2-cyanoethyl)piperidin-4-yloxy)ethyl, 3-(1-(2-cyanoethyl)piperidin-4-yloxy)propyl, 2-(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, (pyrrolidin-2-yl)methyl, 2-(pyrrolidin-1-yl)ethyl, 3-(pyrrolidin-1-yl)propyl, (2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, 5(R)-(2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, (5S)-(2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(N-(2-methoxyethyl)-N-methylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-(2-methoxyethylamino)propyl, 3-(N-(2-methoxyethyl)-N-methylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-(1,2,3-triazol-1-yl)ethyl, 2-(1,2,3-triazol-2-yl)ethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-(1,2,4-triazol-4-yl)ethyl, 4-pyridylmethyl, 2-(4-pyridyl)ethyl, 3-(4-

pyridyl)propyl, 2-(4-pyridyloxy)ethyl, 2-(4-pyridylamino)ethyl, 2-(4-oxo-1,4-dihydro-1-pyridyl)ethyl, 2-(2-oxo-imidazolidin-1-yl)ethyl, 3-(2-oxo-imidazolidin-1-yl)propyl, 2-thiomorpholinoethyl, 3-thiomorpholinopropyl, 2-(1,1-dioxothiomorpholino)ethyl, 3-(1,1-dioxothiomorpholino)propyl, 2-(2-methoxyethoxy)ethyl, 2-(4-methylpiperazin-1-yl)ethyl, 3-(4-methylpiperazin-1-yl)propyl, 3-(methylsulphinyl)propyl, 3-(methylsulphonyl)propyl, 3-(ethylsulphinyl)propyl, 3-(ethylsulphonyl)propyl, 2-(5-methyl-1,2,4-triazol-1-yl)ethyl, morpholino, 2-((N-(3-morpholinopropylsulphonyl)-N-methyl)amino)ethyl, 2-((N-methyl-N-4-pyridyl)amino)ethyl, 3-(4-oxidomorpholino)propyl, 2-(2-(4-methylpiperazin-1-yl)ethoxy)ethyl, 3-(2-(4-methylpiperazin-1-yl)ethoxy)propyl, 2-(2-morpholinoethoxy)ethyl, 3-(2-morpholinoethoxy)propyl, 2-(tetrahydropyran-4-yloxy)ethyl, 3-(tetrahydropyran-4-yloxy)propyl, 2-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)vinyl, 3-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)prop-2-en-1-yl, 1-(2-pyrrolidinylethyl)piperidin-4-ylmethyl, 1-(3-pyrrolidinylpropyl)piperidin-4-ylmethyl, 1-(2-piperidinylethyl)piperidin-4-ylmethyl, 1-(3-piperidinylpropyl)piperidin-4-ylmethyl, 1-(2-morpholinoethyl)piperidin-4-ylmethyl, 1-(3-morpholinopropyl)piperidin-4-ylmethyl, 1-(2-thiomorpholinoethyl)piperidin-4-ylmethyl, 1-(3-thiomorpholinopropyl)piperidin-4-ylmethyl, 1-(2-azetidinyethyl)piperidin-4-ylmethyl, [or] 1-(3-azetidinypropyl)piperidin-4-ylmethyl, 3-morpholino-2-hydroxypropyl, (2*R*)-3-morpholino-2-hydroxypropyl, (2*S*)-3-morpholino-2-hydroxypropyl, 3-piperidino-2-hydroxypropyl, (2*R*)-3-piperidino-2-hydroxypropyl, (2*S*)-3-piperidino-2-hydroxypropyl, 3-pyrrolidin-1-yl-2-hydroxypropyl, (2*R*)-3-pyrrolidin-1-yl-2-hydroxypropyl, (2*S*)-3-pyrrolidin-1-yl-2-hydroxypropyl, 3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, (2*R*)-3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, (2*S*)-3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, 3-(N,N-diethylamino)-2-hydroxypropyl, (2*R*)-3-(N,N-diethylamino)-2-hydroxypropyl, (2*S*)-3-(N,N-diethylamino)-2-hydroxypropyl, 3-(isopropylamino)-2-hydroxypropyl, (2*R*)-3-(isopropylamino)-2-hydroxypropyl, (2*S*)-3-(isopropylamino)-2-hydroxypropyl, 3-(N,N-diisopropylamino)-2-hydroxypropyl, (2*R*)-3-(N,N-diisopropylamino)-2-hydroxypropyl or (2*S*)-3-(N,N-diisopropylamino)-2-hydroxypropyl].

\* \* \* \* \*

The paragraph extending from page 35, line 4 to page 37, line 11 has been amended as follows:

In another aspect R<sup>2</sup> represents ethoxy, trifluoromethoxy, 2,2,2-trifluoroethoxy, 2-hydroxyethoxy, 3-hydroxypropoxy, 2-methoxyethoxy, 3-methoxypropoxy, 2-(methylsulphinyl)ethoxy, 2-(methylsulphonyl)ethoxy, 2-(ethylsulphinyl)ethoxy, 2-(ethylsulphonyl)ethoxy, 2-(N,N-dimethylsulphamoyl)ethoxy, 2-(N-methylsulphamoyl)ethoxy, 2-sulphamoylethoxy, 2-(methylamino)ethoxy, 3-(methylamino)propoxy, 2-(ethylamino)ethoxy, 3-(ethylamino)propoxy, 2-(N,N-dimethylamino)ethoxy, 3-(N,N-dimethylamino)propoxy, 2-(N,N-diethylamino)ethoxy, 3-(N,N-diethylamino)propoxy, 2-(N-methyl-N-methylsulphonylamino)ethoxy, 3-(N-methyl-N-methylsulphonylamino)propoxy, 2-morpholinoethoxy, 3-morpholinopropoxy, 2-piperidinoethoxy, 3-piperidinopropoxy, 2-(methylpiperidino)ethoxy, 3-(methylpiperidino)propoxy, 2-(ethylpiperidino)ethoxy, 3-(ethylpiperidino)propoxy, 2-((2-methoxyethyl)piperidino)ethoxy, 3-((2-methoxyethyl)piperidino)propoxy, 2-((2-methylsulphonyl)ethylpiperidino)ethoxy, 3-((2-methylsulphonyl)ethylpiperidino)propoxy, piperidin-3-ylmethoxy, piperidin-4-ylmethoxy, 2-(piperidin-3-yl)ethoxy, 2-(piperidin-4-yl)ethoxy, 3-(piperidin-3-yl)propoxy, 3-(piperidin-4-yl)propoxy, 2-(piperidin-2-yl)ethoxy, 3-(piperidin-2-yl)propoxy, (1-methylpiperidin-3-yl)methoxy, (1-methylpiperidin-4-yl)methoxy, (1-cyanomethylpiperidin-3-yl)methoxy, (1-cyanomethylpiperidin-4-yl)methoxy, 2-(methylpiperidin-3-yl)ethoxy, 2-(methylpiperidin-4-yl)ethoxy, 2-(1-cyanomethylpiperidin-3-yl)ethoxy, 2-(1-cyanomethylpiperidin-4-yl)ethoxy, 3-(methylpiperidin-3-yl)propoxy, 3-(methylpiperidin-4-yl)propoxy, 3-(1-cyanomethylpiperidin-3-yl)propoxy, 3-(1-cyanomethylpiperidin-4-yl)propoxy, 2-(ethylpiperidin-3-yl)ethoxy, 2-(ethylpiperidin-4-yl)ethoxy, 3-(ethylpiperidin-3-yl)propoxy, 3-(ethylpiperidin-4-yl)propoxy, ((2-methoxyethyl)piperidin-3-yl)methoxy, ((2-methoxyethyl)piperidin-4-yl)methoxy, 2-((2-methoxyethyl)piperidin-3-yl)ethoxy, 2-((2-methoxyethyl)piperidin-4-yl)ethoxy, 3-((2-methoxyethyl)piperidin-3-yl)propoxy, 3-((2-methoxyethyl)piperidin-4-yl)propoxy, (1-(2-methylsulphonylethyl)piperidin-3-yl)methoxy, (1-(2-methylsulphonylethyl)piperidin-4-yl)methoxy, 2-((2-methylsulphonylethyl)piperidin-3-yl)ethoxy, 2-((2-methylsulphonylethyl)piperidin-4-yl)ethoxy, 3-((2-methylsulphonylethyl)piperidin-3-

yl)propoxy, 3-((2-methylsulphonyl)ethyl)piperidin-4-yl)propoxy, 1-isopropylpiperidin-2-ylmethoxy, 1-isopropylpiperidin-3-ylmethoxy, 1-isopropylpiperidin-4-ylmethoxy, 2-(1-isopropylpiperidin-2-yl)ethoxy, 2-(1-isopropylpiperidin-3-yl)ethoxy, 2-(1-isopropylpiperidin-4-yl)ethoxy, 3-(1-isopropylpiperidin-2-yl)propoxy, 3-(1-isopropylpiperidin-3-yl)propoxy, 3-(1-isopropylpiperidin-4-yl)propoxy, 2-(piperidin-4-yloxy)ethoxy, 3-(piperidin-4-yloxy)propoxy, 2-(1-(cyanomethyl)piperidin-4-yloxy)ethoxy, 3-(1-(cyanomethyl)piperidin-4-yloxy)propoxy, 2-(1-(2-cyanoethyl)piperidin-4-yloxy)ethoxy, 3-(1-(2-cyanoethyl)piperidin-4-yloxy)propoxy, 2-(piperazin-1-yl)ethoxy, 3-(piperazin-1-yl)propoxy, (pyrrolidin-2-yl)methoxy, 2-(pyrrolidin-1-yl)ethoxy, 3-(pyrrolidin-1-yl)propoxy, (2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methoxy, 5(R)-(2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methoxy, (5S)-(2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methoxy, (1,3-dioxolan-2-yl)methoxy, 2-(1,3-dioxolan-2-yl)ethoxy, 2-(2-methoxyethylamino)ethoxy, 2-(N-(2-methoxyethyl)-N-methylamino)ethoxy, 2-(2-hydroxyethylamino)ethoxy, 3-(2-methoxyethylamino)propoxy, 3-(N-(2-methoxyethyl)-N-methylamino)propoxy, 3-(2-hydroxyethylamino)propoxy, 2-(1,2,3-triazol-1-yl)ethoxy, 2-(1,2,3-triazol-2-yl)ethoxy, 2-(1,2,4-triazol-1-yl)ethoxy, 2-(1,2,4-triazol-4-yl)ethoxy, 4-pyridylmethoxy, 2-(4-pyridyl)ethoxy, 3-(4-pyridyl)propoxy, 2-(4-pyridyloxy)ethoxy, 2-(4-pyridylamino)ethoxy, 2-(4-oxo-1,4-dihydro-1-pyridyl)ethoxy, 2-(2-oxo-imidazolidin-1-yl)ethoxy, 3-(2-oxo-imidazolidin-1-yl)propoxy, 2-thiomorpholinoethoxy, 3-thiomorpholinopropoxy, 2-(1,1-dioxothiomorpholino)ethoxy, 3-(1,1-dioxothiomorpholino)propoxy, 2-(2-methoxyethoxy)ethoxy, 2-(4-methylpiperazin-1-yl)ethoxy, 3-(4-methylpiperazin-1-yl)propoxy, 3-(methylsulphinyl)propoxy, 3-(methylsulphonyl)propoxy, 3-(ethylsulphinyl)propoxy, 3-(ethylsulphonyl)propoxy, 2-(5-methyl-1,2,4-triazol-1-yl)ethoxy, 2-((N-(3-morpholinopropylsulphonyl)-N-methyl)amino)ethoxy, 2-((N-methyl-N-4-pyridyl)amino)ethoxy, 3-(4-oxidomorpholino)propoxy, 2-(2-(4-methylpiperazin-1-yl)ethoxy)ethoxy, 3-(2-(4-methylpiperazin-1-yl)ethoxy)propoxy, 2-(2-morpholinoethoxy)ethoxy, 3-(2-morpholinoethoxy)propoxy, 2-(tetrahydropyran-4-yloxy)ethoxy, 3-(tetrahydropyran-4-yloxy)propoxy, 2-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)vinyl, 3-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)prop-2-en-1-yloxy, 1-(2-pyrrolidinylethyl)piperidin-4-ylmethoxy, 1-(3-pyrrolidinylpropyl)piperidin-4-ylmethoxy, 1-(2-piperidinylethyl)piperidin-4-ylmethoxy, 1-

(3-piperidinylpropyl)piperidin-4-ylmethoxy, 1-(2-morpholinoethyl)piperidin-4-ylmethoxy, 1-(3-morpholinopropyl)piperidin-4-ylmethoxy, 1-(2-thiomorpholinoethyl)piperidin-4-ylmethoxy, 1-(3-thiomorpholinopropyl)piperidin-4-ylmethoxy, 1-(2-azetidinylethyl)piperidin-4-ylmethoxy, [or] 1-(3-azetidinylpropyl)piperidin-4-ylmethoxy, 3-morpholino-2-hydroxypropoxy, (2R)-3-morpholino-2-hydroxypropoxy, (2S)-3-morpholino-2-hydroxypropoxy, 3-piperidino-2-hydroxypropoxy, (2R)-3-piperidino-2-hydroxypropoxy, (2S)-3-piperidino-2-hydroxypropoxy, 3-pyrrolidin-1-yl-2-hydroxypropoxy, (2R)-3-pyrrolidin-1-yl-2-hydroxypropoxy, (2S)-3-pyrrolidin-1-yl-2-hydroxypropoxy, 3-(1-methylpiperazin-4-yl)-2-hydroxypropoxy, (2R)-3-(1-methylpiperazin-4-yl)-2-hydroxypropoxy, (2S)-3-(1-methylpiperazin-4-yl)-2-hydroxypropoxy, 3-(N,N-diethylamino)-2-hydroxypropoxy, (2R)-3-(N,N-diethylamino)-2-hydroxypropoxy, (2S)-3-(N,N-diethylamino)-2-hydroxypropoxy, 3-(isopropylamino)-2-hydroxypropoxy, (2R)-3-(isopropylamino)-2-hydroxypropoxy, (2S)-3-(isopropylamino)-2-hydroxypropoxy, 3-(N,N-diisopropylamino)-2-hydroxypropoxy, (2R)-3-(N,N-diisopropylamino)-2-hydroxypropoxy or (2S)-3-(N,N-diisopropylamino)-2-hydroxypropoxy.

\* \* \* \* \*

The subparagraph numbered 12) at page 38, line 13, has been amended as follows:


12) C<sub>1-5</sub>alkylX<sup>6</sup>R<sup>29</sup> [C<sub>1-5</sub>alkylX<sup>6</sup>X<sup>29</sup>] (wherein X<sup>6</sup> and R<sup>29</sup> are as defined hereinbefore);

\* \* \* \* \*

The paragraph extending from page 47, line 26 to page 48, line 28 has been amended as follows:

In another aspect of the present invention there is provided the use of compounds of the formula Ia:





(Ia)

[wherein:

ring C,  $R^1$ ,  $R^2$ ,  $n$  and  $Z$  are as defined hereinbefore with the provisos that  $R^2$  is not hydrogen and that  $Z$  is not  $CH_2$  or a direct bond; and

$R^{2a}$  represents hydrogen, halogeno,  $C_{1-3}$ alkyl, trifluoromethyl,  $C_{1-3}$ alkoxy,  $C_{1-3}$ alkylsulphanyl,  $-NR^{3a}R^{4a}$  (wherein  $R^{3a}$  and  $R^{4a}$ , which may be the same or different, each represents hydrogen or  $C_{1-3}$ alkyl), or  $R^{5a}(CH_2)_zX^{1a}$  (wherein  $R^{5a}$  is a [5]4- or 6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano,  $C_{1-4}$ cyanoalkyl,  $C_{1-4}$ alkyl,  $C_{1-4}$ hydroxyalkyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkoxy $C_{1-4}$ alkyl,  $C_{1-4}$ alkylsulphonyl $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxycarbonyl,  $C_{1-4}$ aminoalkyl,  $C_{1-4}$ alkylamino, di( $C_{1-4}$ alkyl)amino,  $C_{1-4}$ alkylamino $C_{1-4}$ alkyl, di( $C_{1-4}$ alkyl)amino $C_{1-4}$ alkyl,  $C_{1-4}$ alkylamino $C_{1-4}$ alkoxy, di( $C_{1-4}$ alkyl)amino $C_{1-4}$ alkoxy and a group  $-(O)_f(C_{1-4}alkyl)_g$ ringD (wherein  $f$  is 0 or 1,  $g$  is 0 or 1 and ring D is a [5]4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from  $C_{1-4}$ alkyl),  $z$  is an integer from 0 to 4 and  $X^{1a}$  represents a direct bond,  $-O-$ ,  $-CH_2-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-NR^{6a}C(O)-$ ,  $-C(O)NR^{7a}-$ ,  $-SO_2NR^{8a}-$ ,  $-NR^{9a}SO_2-$  or  $-NR^{10a}-$  (wherein  $R^{6a}$ ,  $R^{7a}$ ,  $R^{8a}$ ,  $R^{9a}$  and  $R^{10a}$  each independently represents hydrogen,  $C_{1-3}$ alkyl or  $C_{1-3}$ alkoxy $C_{2-3}$ alkyl));

and salts thereof, and prodrugs thereof for example esters and amides, in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in warm-blooded animals such as humans.

\* \* \* \* \*

The first paragraph of Example 15 at page 111, extending from line 9 to line 13, has been amended as follows:

Using a procedure analogous to that described for Example 9, 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (0.13g, 0.4~~mmol~~ [mol]), (prepared as described for the starting material in Example 9), was reacted with 7-hydroxy-4-methylquinoline (80mg, 0.5mol), (Chem. Berich. 1967, 100, 2077), to give **6-methoxy-4-(4-methylquinolin-7-yloxy)-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline** (155mg, 87%).

\* \* \* \* \*

The first paragraph of Example 16 at page 111, extending from line 20 to line 25, has been amended as follows:

Using a procedure analogous to that described for Example 9, 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (0.13g, 0.4~~mmol~~ [mol]), (prepared as described for the starting material in Example 9), was reacted with 2,2,4-trimethyl-1,2-dihydroquinolin-6-ol (95mg, 0.5mmol), (IZV. ACAD. NAVK. SSSR. Ser. Khim. 1981, 9, 2008), to give **6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)-4-(2,2,4-trimethyl-1,2-dihydroquinolin-6-yloxy)quinazoline** (90mg, 47%).

\* \* \* \* \*

The first paragraph of Example 18 at page 112, extending from line 12 to line 16, has been amended as follows:

Using a procedure analogous to that described for Example 9, 4-chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (0.13g, 0.4~~mmol~~ [mol]), (prepared as described for the starting material in Example 10), was reacted with 2,4-dimethyl-7-hydroxyquinoline (87mg, 0.5mmol), (Chem. Berichte, 1903, 36, 4016), to give **4-(2,4-dimethylquinolin-7-yloxy)-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline** (61mg, 33%).

\* \* \* \* \*

The first paragraph of Example 19 at page 112, extending from line 23 to line 27, has been amended as follows:

Using a procedure analogous to that described for Example 9, 4-chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (0.13g, 0.4~~mmol~~ [mol]), (prepared as described for the starting material in Example 10), was reacted with 6-hydroxy-2*H*-4*H*-1,4-benzoxazin-3-one (83mg, 0.5~~mmol~~ [mol]), (J. Chem. Soc. C, 1971, 2696), to give 6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(3-oxo-2*H*-4*H*-1,4-benzoxazin-6-yloxy)quinazoline (158mg, 88%).

\* \* \* \* \*

The first paragraph of Example 20 at page 113, extending from line 2 to line 6, has been amended as follows:

Using a procedure analogous to that described for Example 9, 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (0.13g, 0.4~~mmol~~ [mol]), (prepared as described for the starting material in Example 9), was reacted with 6-hydroxy-2*H*-4*H*-1,4-benzoxazin-3-one (83mg, 0.5~~mmol~~ [mol]), (J. Chem. Soc. C, 1971, 2696), to give 6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)-4-(3-oxo-2*H*-4*H*-1,4-benzoxazin-6-yloxy)quinazoline (170mg, 94%).

\* \* \* \* \*

The first paragraph of Example 71 at page 140, extending from line 12 to line 23, has been amended as follows:

Methoxyacetaldehyde (368mg, 3.47 ~~mmol~~ [mol]) (freshly distilled) followed by sodium triacetoxyborohydride (552mg, 2.6mol) were added to a solution of 6-methoxy-4-(2-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline (726mg, 1.74mmol), (prepared as described in Example 70), in a mixture of methylene chloride (15ml) and methanol (15ml). After stirring for 1.5 hours at ambient temperature, saturated sodium hydrogen carbonate was added. The volatiles were removed under vacuum and the residue was partitioned between methylene chloride and water. The organic layer, was separated, washed with water, brine, dried (MgSO<sub>4</sub>) and the volatiles were removed by evaporation. The residue was purified by column chromatography eluting with methylene chloride/methanol

(80/20). After removal of the solvent, the residue was triturated with ether, collected by filtration, washed with ether and dried under vacuum at 60°C to give **6-methoxy-7-(1-(2-methoxyethyl)piperidin-4-ylmethoxy)-4-(2-methylindol-5-yloxy)quinazoline** (392mg, 47%).

\* \* \* \* \*

The first paragraph of Example 90 at page 153, extending from line 4 to line 12, has been amended as follows:

A solution of 7-hydroxy-4-(2-methylindol-5-yloxy)quinazoline (423 mg, 1.45 mmol [mol]), (prepared as described for the starting material in Example 82), triphenylphosphine (685 mg, 2.61 mmol), 4-hydroxymethyl-1-*tert*-butoxycarbonylpiperidine (500 mg, 2.32 mmol), (prepared as described for the starting material in Example 10), and diisopropyl azodicarboxylate (528 mg, 2.61 mmol) in methylene chloride (18 ml) was stirred overnight at ambient temperature. The mixture was then poured onto a column of silica and eluted with ethyl acetate. After evaporation of the solvent, the residue was triturated with ether, filtered, and dried under vacuum to give **7-(1-*tert*-butoxycarbonylpiperidin-4-ylmethoxy)-4-(2-methylindol-5-yloxy)quinazoline** (478 mg, 68 %).

\* \* \* \* \*

The paragraph of Example 126 at page 168 extending from line 19 to line 24, has been amended as follows:

A solution of 2-(piperidin-4-yl)-1-ethanol (830 mg, 6.4 mmol) in DMF (5 ml) containing *tert*butyl dicarbonate anhydride (1.4 g, 6.4 mmol [mol]) was stirred at ambient temperature for 48 hours. After removal of the volatiles under vacuum, the residue was partitioned between ether and water. The organic layer was separated, washed with water, brine, dried (MgSO<sub>4</sub>) and evaporated to give **4-(2-hydroxyethyl)-(1-*tert*-butoxycarbonyl)piperidine** (1 g, 68 %).

\* \* \* \* \*

The first paragraph of Example 157 at page 189, extending from line 20 to line 28, has been amended as follows:

2-Chloro-3-fluoro-7-methoxyquinoline (210~~mg~~ [g], 1mmol), (prepared as described for the starting material in Example 157), in anhydrous THF (1ml) was added to a mixture of copper(I)bromide (570mg, 4.0mmol) and methylmagnesium bromide (3.0M solution in diethyl ether, 2.7ml, 8mmol) in anhydrous THF (20ml) at -78°C. The mixture was stirred for 1 hour at -78°C, allowed to warm to ambient temperature and then stirred for a further 18 hours. Saturated aqueous ammonium chloride solution and 5N aqueous sodium hydroxide solution (pH 12) were added and the product extracted with ethyl acetate (3x). The organic solution was washed with water, brine, dried (MgSO<sub>4</sub>) and evaporated to dryness to yield 3-fluoro-7-methoxy-2-methylquinoline (0.17g, 91%).

\* \* \* \* \*

The table in Example 192 at page 218, extending from line 1 to line 2, has been amended as follows:

Elemental analysis:	Found	C 66.4	H 6.9	N 12.4
[C <sub>2</sub> H <sub>1</sub> N <sub>1</sub> O <sub>1</sub> ; 1 HCl, 2 H <sub>2</sub> O]				
<u>C<sub>25</sub>H<sub>30</sub>N<sub>4</sub>O<sub>4</sub></u>	Requires	C 66.7	H 6.7	N 12.4%

**IN THE CLAIMS:**

Claims 1-30 have been cancelled and replaced by new claims 31-60.